

ON ROTATIONALLY INVARIANT SHRINKING GRADIENT RICCI SOLITONS

BRETT KOTSCHWAR

ABSTRACT. In this paper we study the gradient Ricci shrinking soliton equation on rotationally symmetric manifolds of dimension three and higher and prove that the only complete examples of such metrics on S^n , $\mathbb{R}^n$ and $\mathbb{R} \times S^{n-1}$ are, respectively, the round, flat, and standard cylindrical metrics.

1. INTRODUCTION

Recall that a Riemannian manifold $(\mathcal{M}^{n+1}, g)$ is said to be a *shrinking gradient Ricci soliton* provided

$$(1) \quad Rc(g) + \nabla \nabla f - \lambda g = 0$$

for some $f \in C^\infty(\mathcal{M}^{n+1})$ and $\lambda > 0$. The analogous objects in the cases $\lambda = 0$ and $\lambda < 0$ are known as *steady* and *expanding* solitons, respectively, and the triple (g, f, λ) is referred to as a *(gradient) soliton structure* on the manifold $\mathcal{M}^{n+1}$. Solitons correspond to self-similar solutions of the Ricci flow and arise commonly in the analysis of its singularities.

After the class of Einstein metrics, perhaps the most natural place to look for solitons is among the rotationally invariant metrics, and, in all dimensions greater than one, this class has been shown to contain complete, non-trivial examples of steady and expanding solitons (cf., e.g., [I] and [RF2]). However, one does not expect to find corresponding examples in the shrinking case. The purpose of the present work is to confirm this expectation. Our main result is the following

Theorem 1. *Suppose $n \geq 2$, and (g, f, λ) is a complete, rotationally invariant shrinking soliton structure on a manifold $\mathcal{M}^{n+1}$ diffeomorphic to one of S^{n+1} , $\mathbb{R}^{n+1}$ or $\mathbb{R} \times S^n$. Then,*

- (1) *If $\mathcal{M}^{n+1} \approx S^{n+1}$, g is isometric to a round sphere and $f \equiv \text{const}$.*
- (2) *If $\mathcal{M}^{n+1} \approx \mathbb{R}^{n+1}$, g is flat.*
- (3) *If $\mathcal{M}^{n+1} \approx \mathbb{R} \times S^n$, g is isometric to the standard cylinder $dr^2 + \omega_0^2 g_{S^n}$ of radius $\omega_0 = \sqrt{(n-1)/\lambda}$ and $f = f(r) = (n-1)r^2/(2\omega_0^2) + \text{linear}$.*

This theorem is known (in greater generality) in dimensions less than four. In dimension two, work of Hamilton ([H], [H2]) shows that the only complete shrinking solitons are the flat metric on $\mathbb{R}^2$ and the round metric on S^2 . In fact, as Hamilton observes in [H], surface solitons are necessarily rotationally symmetric—an observation used, for example, by Chen, Lu, and Tian in their note [CLT] to provide an alternative proof of the uniqueness of the constant curvature soliton on S^2 . In a sense, then, our result is an extension of these two-dimensional findings to higher dimensions.

In dimension three, gradient solitons need not be rotationally symmetric, however Perelman [P2] has shown that the only complete examples of shrinking solitons with non-negative sectional curvature are the flat metric on $\mathbb{R}^3$, the round metric on S^3 , and the standard metric on the cylinder $\mathbb{R} \times S^2$. As the Hamilton-Ivey estimate ([H], [I]) implies that three-dimensional ancient solutions are necessarily of non-negative curvature, Perelman's argument classifies all complete three-dimensional shrinking solitons

In higher dimensions, a consequence of the recent work of Böhm and Wilking [BW], is that the only compact shrinking soliton with 2-positive curvature operator is the round sphere. In the Kähler category, Feldman, Ilmanen, and Knopf [FIK] have obtained results similar to our own for Kähler-Ricci shrinking solitons under the assumption of $U(n)$ -invariance. In particular, their Proposition 9.2– that the flat metric is the only complete $U(n)$ -invariant gradient shrinking soliton on $\mathbb{C}^n$ –is the Kähler analog of the second case of our Theorem 1. Their paper also provides non-trivial examples of $U(n)$ -invariant gradient solitons (of all types) on other spaces.

To prove Theorem 1, we show first that, under the constraint (1), the rotational symmetry of a non-flat metric implies the rotational symmetry of the gradient function. This reduces the proof of the proposition to the study of a certain second order system of non-linear ODEs. By a change of variables due to Robert Bryant and Tom Ivey [RF2], we are able to further reduce the problem to the study of an equivalent first-order system amenable to phase-plane analysis. For the case $\mathcal{M}^{n+1} \approx S^{n+1}$, we show that any candidate metric must have positive curvature operator and then apply the result of Böhm and Wilking. It is no doubt possible to prove this case solely from the analysis of the system of ODE (without using any of the dynamic properties of the Ricci flow), however, since we feel the result may not come as a surprise to experts, we content ourselves here with a proof by the most readily available means. For the non-compact cases, we use the criterion of completeness to eliminate all but the two standard metrics by their asymptotic behavior.

2. THE SOLITON CONDITION FOR ROTATIONALLY SYMMETRIC MANIFOLDS.

2.1. The warped-product metric. Fix $n > 1$ and let $\tilde{g}$ denote the metric on S^n of constant sectional curvature 1. For $-\infty \leq A < \Omega \leq \infty$, and positive functions $\omega \in C^\infty(A, \Omega)$, we shall consider the warped-product metric $g = dr^2 + \omega^2(r)\tilde{g}$ on the cylinder $\mathcal{C}_{A,\Omega} := (A, \Omega) \times S^n$. For this metric, we have the following standard

Lemma 2. *The metric $g = dr^2 + \omega(r)^2\tilde{g}$ on $\mathcal{C}_{0,\Omega}$ extends to a smooth metric on $B_\Omega(\mathbf{0}) \subset \mathbb{R}^{n+1}$ if and only if $\Omega > 0$ and*

$$\lim_{r \rightarrow 0} \omega(r) = 0, \quad \lim_{r \rightarrow 0} \omega'(r) = 1, \quad \text{and} \quad \lim_{r \rightarrow 0} \frac{d^{2k}\omega}{dr^{2k}}(r) = 0 \quad \text{for all } k.$$

The metric extends to a smooth metric on S^{n+1} if and only if in addition $\Omega < \infty$ and

$$\lim_{r \rightarrow \Omega} \omega(r) = 0, \quad \lim_{r \rightarrow \Omega} \omega'(r) = -1, \quad \text{and} \quad \lim_{r \rightarrow \Omega} \frac{d^{2k}\omega}{dr^{2k}}(r) = 0 \quad \text{for all } k.$$

Proof. See, e.g., [CK], Lemma 2.10. □

Thus we may accomplish the proof of Theorem 1 by the study of soliton structures on $\mathcal{C}_{A,\Omega}$. We begin by recording the expressions of a few geometric quantities associated to g .

The sectional curvatures of g are

$$(2) \quad \nu_1 = -\frac{\omega''}{\omega} \quad \text{and} \quad \nu_2 = \frac{1 - (\omega')^2}{\omega^2},$$

where ν_1 denotes the curvature of two-planes tangent to the radial direction, and ν_2 the curvature of orbital two-planes. The Ricci curvature of g has the form

$$(3) \quad Rc(g) = -n \frac{\omega''}{\omega} dr^2 + ((n-1)(1 - (\omega')^2) - \omega\omega'') \tilde{g}$$

where the prime denotes differentiation with respect to r .

A routine computation also shows that for a smooth function f on $\mathcal{C}_{A,\Omega}$, one has

$$\nabla \nabla f = \begin{cases} \nabla_0 \nabla_0 f &= f_{00} \\ \nabla_0 \nabla_i f &= f_{i0} - (\omega'/\omega) f_i \\ \nabla_i \nabla_j f &= \tilde{\nabla}_i \tilde{\nabla}_j f + \omega\omega' f_0 \tilde{g}_{ij} \end{cases}$$

in local coordinates $(\theta^0 = r, \theta^i)$.¹

In these coordinates, (1) has the expression

$$(4) \quad f_{00} - n \frac{\omega''}{\omega} - \lambda = 0$$

$$(5) \quad f_{0i} - \frac{\omega'}{\omega} f_i = 0$$

$$(6) \quad \tilde{\nabla}_i \tilde{\nabla}_j f + [(n-1)(1 - (\omega')^2) - \omega\omega'' - \omega\omega' f_0 - \lambda\omega^2] \tilde{g}_{ij} = 0.$$

The above system involves the partial derivatives of f , and despite the rotational symmetry of the metric and the Ricci curvature, there is no *a priori* reason to assume that f shares this symmetry. However, as we show next, this symmetry is implied unless g is flat, and thus for the proof of Theorem 1, there is no loss in restricting our attention to the case $f = f(r)$.

When f is a radial function, equations (4)-(6) reduce to

$$(7) \quad \begin{cases} f'' - \lambda &= n\omega''/\omega \\ \omega\omega' f' - \lambda\omega^2 &= \omega\omega'' + (n-1)((\omega')^2 - 1). \end{cases}$$

Proposition 3. *Suppose that (g, f, λ) is a complete, rotationally symmetric gradient shrinking soliton structure on $\mathcal{M}^{n+1} \approx S^{n+1}$, $\mathbb{R}^{n+1}$, or $\mathbb{R} \times S^n$.*

If the function f is not rotationally symmetric, then $\mathcal{M}^{n+1} \cong \mathbb{R}^{n+1}$ and g is the flat metric.

Proof. Write $g = dr^2 + \omega^2(r)\tilde{g}$ for $r \in (A, \Omega)$, and fix local coordinates $r = \theta^0, \theta^1, \dots, \theta^n$ on a neighborhood $(A, \Omega) \times \mathcal{U}$ about any point. Observe that for each fixed r , equation (6) is a tensorial identity on S^n , and that we may therefore differentiate it using the Levi-Civita connection $\tilde{\nabla}$ of $\tilde{g}$ to obtain

$$\tilde{\nabla}_k \tilde{\nabla}_i \tilde{\nabla}_j f + \omega\omega' f_{0k} \tilde{g}_{ij} = 0.$$

¹ Here and throughout, when working in coordinates $\theta^0, \dots, \theta^n$, we shall use roman letters to denote indices $1, \dots, n$ and use a tilde to denote quantities (Levi-Civita connection, curvature, etc.) associated to the metric $\tilde{g}$ on S^n . In particular, $\tilde{\nabla}_i \tilde{\nabla}_j f$ represents the hessian of $f(r, \cdot)$ considered as a function on S^n .

Hence

$$\tilde{\nabla}_k \tilde{\nabla}_i \tilde{\nabla}_j f - \tilde{\nabla}_i \tilde{\nabla}_k \tilde{\nabla}_j f = \omega \omega' (f_{0i} \tilde{g}_{jk} - f_{0k} \tilde{g}_{ij}).$$

On the other hand, since $\tilde{R}_{ijkl} = \tilde{g}_{il} \tilde{g}_{jk} - \tilde{g}_{ik} \tilde{g}_{jl}$, the standard commutation identities imply

$$\begin{aligned} \tilde{\nabla}_k \tilde{\nabla}_i \tilde{\nabla}_j f - \tilde{\nabla}_i \tilde{\nabla}_k \tilde{\nabla}_j f &= -\tilde{R}_{kijl} \tilde{g}^{lm} f_m \\ &= -(\tilde{g}_{ij} \tilde{g}_{kl} - \tilde{g}_{il} \tilde{g}_{jk}) \tilde{g}^{lm} f_m \\ &= f_i \tilde{g}_{jk} - f_k \tilde{g}_{ij}. \end{aligned}$$

Combining the two, we find

$$(\omega \omega' f_{0i} - f_i) \tilde{g}_{jk} = (\omega \omega' f_{0k} - f_k) \tilde{g}_{ij},$$

and tracing yields

$$(n-1)(\omega \omega' f_{0i} - f_i) = 0$$

for all $i = 1, \dots, n$. Together with (5), we conclude

$$(8) \quad (n-1) |\tilde{\nabla} f|_{\tilde{g}}^2 [1 - (\omega')^2] = 0.$$

Since we assume $n > 1$, if $(Xf)(r_0, \theta_0) \neq 0$ for some $X \in T_{(r_0, \theta_0)} \mathcal{M}^{n+1}$ tangent to the S^n factor, we must have $|\omega'| \equiv 1$ and $\omega'' \equiv 0$ on an interval $(a, b) \subset (A, \Omega)$ containing r_0 . But, by equation (2), this means that $\nu_1 = \nu_2 = 0$ on (a, b) . We claim that $\nu_1 = \nu_2 = 0$ on the entire interval (A, Ω) .

Let

$$\beta = \sup \{r < \Omega \mid (\omega')^2 = 1 \text{ on } (a, r)\}.$$

If $\beta < \Omega$, by equations (6) and (8), we must have that $(\tilde{\nabla} \tilde{\nabla} f)(\beta, \cdot) = 0$, $\omega'(\beta) = \sigma \in \{\pm 1\}$, $\omega''(\beta) = 0$, $\omega(\beta) > 0$, and $f_0(\beta, \cdot) = \sigma \lambda \omega(\beta)$. Moreover, for some small ϵ , f is a function only of r on $[\beta, \beta + \epsilon)$ and on this interval, f' and ω satisfy the system (7), with the above initial conditions. But one may check that the functions

$$\bar{\omega}(r) = \sigma(r - \beta) + \omega(\beta) \quad \text{and} \quad \bar{f}'(r) = \lambda((r - \beta) + \omega(\beta))$$

also satisfy (7) and agree with ω and f at $r = \beta$. Therefore, by uniqueness², these solutions must coincide and it follows that $(\omega')^2 = (\bar{\omega}')^2 = 1$ on the interval $[\beta, \beta + \epsilon)$, contradicting our choice of β . So g is flat on $(a, \Omega) \times S^n$. Using a similar argument at the other endpoint a , can show that g must be flat on the entire cylinder $(A, \Omega) \times S^n$. But this means either $\omega' \equiv 1$ or $\omega' \equiv -1$, so g cannot extend to a smooth metric on the sphere S^{n+1} or to a complete metric on the cylinder. The only possibility is $\omega' \equiv 1$ and $\mathcal{M}^{n+1} \approx \mathbb{R}^{n+1}$. $\square$

²Writing $x = \omega'$, and $u = f'$, we may recast (7) as a first-order system

$$\begin{cases} \omega' &= x := F(\omega, x, u) \\ x' &= xu - \lambda\omega + (n-1)\frac{1-x^2}{\omega} := G(\omega, x, u) \\ u' &= n\frac{xu}{\omega} + (n-1)\left(n\frac{1-x^2}{\omega^2} - \lambda\right) := H(\omega, x, u). \end{cases}$$

Since F, G, H are C^∞ on the region $\{\omega \neq 0\}$, the asserted uniqueness follows from standard ODE theory.

2.2. An equivalent first-order system and its linearization. In view of the result of the last section, we now assume $f = f(r)$. We are interested in solutions $(\omega(r), f(r))$ to the system (7) for which ω is strictly positive.

As Ivey observes in [RF2], (7) is invariant under translations of r and f . By the introduction of the variables

$$x = \omega' \quad \text{and} \quad y = n\omega' - \omega f'$$

which share this invariance, and of an independent variable t which satisfies $dt = 1/\omega dr$, one obtains the first-order system

$$(9) \quad \begin{cases} \frac{d\omega}{dt} &= x\omega \\ \frac{dx}{dt} &= x^2 - xy + n - 1 - \lambda\omega^2 \\ \frac{dy}{dt} &= xy - nx^2 - \lambda\omega^2. \end{cases}$$

Any solution to (7) gives rise to a trajectory of (9) and conversely, from a trajectory $(\omega(t), x(t), y(t))$ of (9), one may recover r , $\omega(r)$, and $f(r)$ by a succession of quadratures (see [I]). Consequently, it suffices to analyze solutions to the simpler system (9). We take as coordinates (ω, x, y) on the phase space $\mathbb{R}^3$ and restrict our attention to trajectories lying in the half space $\omega > 0$.

For $n > 1$, system (9) has two equilibrium points: $P_0 := (0, 1, n)$ and $P_1 := (0, -1, -n)$. Denoting the right hand side of (9) by Φ , one finds

$$d\Phi_{P_0} = -d\Phi_{P_1} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 2-n & -1 \\ 0 & -n & 1 \end{pmatrix}$$

which has eigenvalues 2, 1, and $1 - n$. Since we assume $n \geq 2$, both P_0 and P_1 are saddle points: P_0 (P_1) lying at the intersection of a two-dimensional unstable (stable) manifold and a one-dimensional stable (unstable) manifold. In particular, there is a one-parameter family of trajectories in the half-space $\omega > 0$ initially tangent to $(1, 0, 0)$, among which, in light of Lemma (2), lie the trajectories which give rise to smooth solutions on S^{n+1} and $\mathbb{R}^{n+1}$ (see Examples (5) and (6) below). Trajectories which correspond to smooth solutions on S^{n+1} must, in addition, tend to P_1 as $t \rightarrow \infty$, and hence lie in the intersection of the global unstable and stable manifolds of P_0 and P_1 , respectively.

Remark 4. If $L : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is the map $(\omega, x, y) \mapsto (\omega, -x, -y)$, then from any solution $\gamma(t) = (\omega(t), x(t), y(t))$ of (9) on (S, T) , one may obtain a new solution

$$\bar{\gamma}(t) := L(\gamma(\tau(t))) = (\omega(\tau(t)), -x(\tau(t)), -y(\tau(t)))$$

on an appropriate interval $(\bar{S}, \bar{T})$ where τ is chosen to satisfy $\frac{d\tau}{dt} = -1$ and $\tau(\bar{S}) = T$, $\tau(\bar{T}) = S$.

By use of this device, one immediately obtains that the global stable and unstable manifolds of P_i , S_i and U_i , $i = 1, 2$ are related by $L(S_0) = U_1$, $L(U_0) = S_1$, and moreover, if a set $V \subset \mathbb{R}^3$ is preserved by the system (9) for increasing t (i.e., $\gamma(t_0) \in V$ implies $\gamma(t) \in V$ for $t > t_0$, as long as the solution is defined), then $L(V)$ is preserved by the system for decreasing t .

2.3. The standard examples. At this juncture, it is worthwhile to recall the standard solutions to (7) and locate the corresponding solution to (9) in ωxy -space. The content of Theorem 1 is that this list essentially exhausts the possibilities for complete solutions.

Example 5. (*Round sphere*) The condition $f \equiv \text{const}$ in (7) leads to the constant curvature soliton structure

$$\begin{cases} \omega(r) &= \sqrt{\frac{\pi}{\lambda}} \sin\left(\sqrt{\frac{\lambda}{n}} r\right) \\ f(r) &= \text{const} \end{cases}$$

on the sphere S^{n+1} . The corresponding trajectory in ωxy -space is the elliptical arc $\{nx^2 + \lambda\omega^2 = n\}$ lying in the plane $\{y = nx\}$ joining P_0 and P_1 .

Example 6. (*Gaussian soliton*) The condition $\omega' \equiv 1$ in (7) leads to the flat soliton structure

$$(10) \quad \begin{cases} \omega(r) &= r \\ f(r) &= \frac{\lambda}{2} r^2 + \text{linear} \end{cases}$$

on $\mathbb{R}^{n+1}$ and corresponds to the trajectory $y = n - \omega^2$ in the plane $\{x = 1\}$ emanating from P_0 . Applying the device of Remark (4), one obtains a similar trajectory in the plane $\{x = -1\}$, satisfying $y = -n + \lambda\omega^2$ and tending to P_1 as $t \rightarrow \infty$.

Example 7. (*Standard cylinder*) The condition $\omega' \equiv 0$ in (7) leads to the structure

$$\begin{cases} \omega(r) &= \sqrt{\frac{n-1}{\lambda}} \\ f(r) &= \frac{\lambda}{2} r^2 + \text{linear} \end{cases}$$

on $\mathbb{R} \times S^n$ corresponding to the line $\{(\sqrt{(n-1)/\lambda}, 0, y)\}$ in ωxy -space.

Remark 8. The flat and cylindrical solutions described in the previous two examples are the only for which $\omega' \equiv \text{const}$, and the corresponding trajectories in ωxy -space describe the intersections of the planes $\{x = 1\}$, $\{x = 0\}$, and $\{x = -1\}$ with the set $\{x^2 - xy + n - 1 - \lambda\omega^2 = 0\}$. This leads to an observation which we shall have repeated occasion to use in the sequel: the only solutions $\gamma(t)$ of (9) for which $\frac{dx}{dt}(t_0) = 0$ and $x(t_0) = 1, 0, \text{ or } -1$ at some t_0 are, by uniqueness, those for which $x(t) \equiv 1, 0, \text{ or } -1$, respectively.

Finally we note (as does Ivey in [RF2]) that by taking $\lambda = 0$ in the x and y -components of (9) one recovers the analogous system for rotationally symmetric solutions to the steady soliton equation. Thus the trajectories of (9) which lie in the plane $\{\omega = 0\}$ are naturally associated with steady soliton structures (although, of course, the warping function $\omega(r)$ of these structures no longer corresponds directly to the ω -coordinate). In particular, the (one-dimensional) intersection of the unstable manifold of P_0 with the plane $\{\omega = 0\}$ contains two candidates for a smooth steady soliton on $\mathbb{R}^{n+1}$: one with negative sectional curvature near the origin, which turns out to be incomplete, and one with positive curvature near the origin, which is the well-known *Bryant soliton*—a complete steady soliton on $\mathbb{R}^{n+1}$ of strictly positive curvature.

3. PROOF OF THEOREM 1

In what follows, $\gamma(t) = (\omega(t), x(t), y(t))$ will represent a trajectory of (9) defined for t in what we may take to be a *maximal* interval (S, T) with $-\infty \leq S < T \leq \infty$. To reduce the clutter of our expressions, we shall usually suppress the dependence of the components of the trajectory on the parameter t .

3.1. Some invariant sets. We begin our analysis of trajectories of (9) by observing that the second and third derivatives of the x -component have the following convenient expressions.

Lemma 9. *The x -component of any trajectory $\gamma(t)$ of (9) satisfies*

$$(11) \quad \frac{d^2x}{dt^2} = (n-1)x(x^2-1) + (3x-y)\frac{dx}{dt}$$

and

$$(12) \quad \frac{d^3x}{dt^3} = 2x[(2n-1)x-y]\frac{dx}{dt} + 2\left(\frac{dx}{dt}\right)^2 + (3x-y)\frac{d^2x}{dt^2}.$$

Proof. We compute

$$\begin{aligned} \frac{d^2x}{dt^2} &= (2x-y)\frac{dx}{dt} - x^2y + nx^3 - \lambda\omega^2x \\ &= (n-1)x(x^2-1) + (3x-y)\frac{dx}{dt}, \end{aligned}$$

and

$$\begin{aligned} \frac{d^3x}{dt^3} &= \left[(n-1)(3x^2-1) + \frac{dx}{dt} - \frac{dy}{dt}\right]\frac{dx}{dt} + 2\left(\frac{dx}{dt}\right)^2 + (3x-y)\frac{d^2x}{dt^2} \\ &= 2x[(2n-1)x-y]\frac{dx}{dt} + 2\left(\frac{dx}{dt}\right)^2 + (3x-y)\frac{d^2x}{dt^2}. \end{aligned}$$

□

With the above expressions we may easily establish the following qualitative results on the behavior of trajectories of the system.

Lemma 10. *The regions*

$$\left\{x \geq 1, \frac{dx}{dt} \geq 0\right\}, \left\{x \leq -1, \frac{dx}{dt} \leq 0\right\}, \text{ and } \{y \leq 0\}$$

are preserved under system (9) for increasing t , and

$$\left\{x \geq 1, \frac{dx}{dt} \leq 0\right\}, \left\{x \leq -1, \frac{dx}{dt} \geq 0\right\}, \text{ and } \{y \geq 0\},$$

are preserved for decreasing t .

Proof. By (11), if $\frac{dx}{dt}(t_0) = 0$ at some t_0 with $x(t_0) > 1$, then $\frac{d^2x}{dt^2}(t_0) > 0$, and hence both $\frac{dx}{dt}$ and x continue to increase. By Remark 8, we cannot have $\frac{dx}{dt} = 0$ and $x = 1$ simultaneously unless $x \equiv 1$. The argument for the case $x \leq -1$ and $\frac{dx}{dt} \leq 0$ is similar.

To see that $\{y \leq 0\}$ is preserved, observe that

$$\frac{dy}{dt} = -nx^2 - \lambda\omega^2 < 0$$

whenever $y = 0$.

The preservation of the remaining sets for decreasing t follows by applying the results already obtained to the trajectory $\bar{\gamma}(t) = L(\gamma(\tau))$ constructed as in Remark (4). □

Lemma 11.

- (1) If there exists $t_0 \in (S, T)$ at which $x(t_0) = 0$, $y(t_0) \leq 0$, and $\frac{dx}{dt}(t_0) > 0$, then $x(t)$ (and $\frac{dx}{dt}$) increase until $\gamma(t)$ enters the region $\{x > 1\}$.
- (2) If there exists $t_0 \in (S, T)$ at which $x(t_0) = 0$, $y(t_0) \geq 0$, and $\frac{dx}{dt}(t_0) > 0$, then $x(t) < 0$ and $\frac{dx}{dt}(t) > 0$ for all $t < t_0$ and there exists a $t_1 \leq t_0$ such that $x(t) < -1$ for all $t < t_1$.

In particular, in view of Lemma 10, if the trajectory γ enters the region $\{x < 0\}$, either it remains there or eventually lies in the region $\{x > 1\}$.

Proof. In case (1), we have $y(t) < 0$ for all $t > t_0$ by Lemma 10, and $\frac{d^2x}{dt^2}(t_0) \geq 0$ by equation (11). Since by (12), $\frac{d^3x}{dt^3} > 0$ on the region

$$\left\{ x \geq 0, \frac{dx}{dt} > 0, \frac{d^2x}{dt^2} \geq 0 \right\},$$

we have $\frac{d^2x}{dt^2}(t) > 0$ for all $t > t_0$. Consequently, $x(t) > \frac{dx}{dt}(t_0)(t - t_0)$. As bounds on x imply bounds on the derivatives of ω and y , the solution cannot expire while $0 < x < 1$. Since the interval (S, T) is maximal, $x > 1$ eventually.

Case (2) then follows from case (1) by considering again the trajectory $\bar{\gamma}(t)$ constructed according to the device in Remark 4: If $x(t_0) = 0$, $y(t_0) \geq 0$ and $\frac{dx}{dt}(t_0) > 0$, then $\bar{x}(t_0) = 0$, $\bar{y}(t_0) \leq 0$ and $\frac{d\bar{x}}{dt}(t_0) > 0$, and the corresponding interval of definition $(\bar{S}, \bar{T})$, satisfying $\tau(\bar{T}) = S$, $\tau(\bar{S}) = T$ will also be maximal. $\square$

3.2. Proof of the case $\mathcal{M}^{n+1} \approx S^{n+1}$.

Theorem 12. *Suppose (S^{n+1}, g) is a rotationally symmetric shrinking soliton. Then g has positive curvature operator.*

Proof. Recall that, for a rotationally symmetric metric, the positivity of the curvature operator is implied by that of the sectional curvatures, and these curvatures have the expressions

$$\nu_1 = -\frac{1}{\omega^2} \frac{dx}{dt}, \text{ and } \nu_2 = \frac{(1-x^2)}{\omega^2}$$

in terms of the (ω, x, y) coordinates.

Any trajectory $\gamma(t)$ of system (9) which corresponds to a smooth soliton structure on the sphere must tend to $P_0 = (0, 1, n)$ as $t \rightarrow -\infty$ and to $P_1 = (0, -1, -n)$ as $t \rightarrow \infty$. Thus, by Lemma 10, we must have $-1 < x(t) < 1$ (hence $\nu_2(t) > 0$) for all t and $\frac{dx}{dt} < 0$ at least initially. We wish to show $\frac{dx}{dt} < 0$ for all t .

By equation (11) of Lemma 9, $\frac{d^2x}{dt^2} < 0$ at critical points of x in the region $\{0 < x < 1\}$, so $\frac{dx}{dt}$ remains strictly negative on this region and, by Remark 8, cannot vanish on $\{x = 0\}$. Since $\gamma(t)$ must tend to P_1 as $t \rightarrow \infty$, it must, in particular, enter the region $\{x < 0\}$, and, in view of Lemma 11, remain there for all subsequent t . Since $\frac{d^2x}{dt^2} > 0$ at critical points of x in the region $\{-1 < x < 0\}$, $\frac{dx}{dt}$ must therefore remain strictly negative if x is to approach -1 . So $\frac{dx}{dt} < 0$ always, and thus for any trajectory emanating from P_0 and tending to P_1 we have $\nu_1 > 0$ and $\nu_2 > 0$ for all t as claimed.

Taking limits, one finds that at the ‘‘poles’’ $r = A$ and $r = \Omega$, the sectional curvatures agree and are at least non-negative. One may therefore apply Lemma 8.2 of [H4] to conclude that the curvature operator $Rm(g) : \wedge^2 \rightarrow \wedge^2$ is of constant rank, and therefore strictly positive everywhere. $\square$

That g has constant sectional curvature is then an immediate consequence of the following

Theorem. (*Böhm and Wilking [BW]: Theorem 1*) *On a compact manifold the normalized Ricci flow evolves a Riemannian metric with 2-positive curvature operator to a limit metric with constant sectional curvature.*

That $f \equiv \text{const}$ in case (1) of Proposition 1, then follows by substituting $\nu_1 = \nu_2 = \text{const}$ into (7) or, alternatively, by considering the identity

$$R + |\nabla f|^2 - 2\lambda f = \text{const}$$

valid on any gradient Ricci shrinking soliton (see, e.g., [RF2]). If f attains its maximum and minimum at the points x_M and x_m , respectively, then the identity implies $f(x_M) = f(x_m)$ since R is constant.

3.3. The asymptotic behavior of trajectories corresponding to complete, non-compact metrics. Hereafter, we shall consider solutions $\gamma(t)$ satisfying one or both of the conditions

$$(13) \quad \int_{t_0}^T \omega(\sigma) d\sigma = \infty = \lim_{t \rightarrow T} r(t)$$

and

$$(14) \quad \int_S^{t_0} \omega(\sigma) d\sigma = \infty = -\lim_{t \rightarrow S} r(t)$$

for any $t_0 \in (S, T)$. Condition (13) is necessarily satisfied by any trajectory corresponding to a complete metric on $\mathbb{R}^{n+1}$ and both (13) and (14) are necessarily satisfied by any trajectory corresponding to a complete metric on $\mathbb{R} \times S^n$. As we shall see, these conditions impose rather stringent conditions on the asymptotic behavior of a trajectory.

We remark that if $\gamma(t)$ satisfies (13) then $\bar{\gamma}(t) = L(\gamma(\tau))$ satisfies (14) and vice-versa. Thus, from the following results, which apply to trajectories satisfying the “forwards” condition (13), we may easily obtain corresponding results for trajectories satisfying the “backwards” condition (14). These results will be collected in Lemma 17, below.

We begin with a simple consequence of the forwards extendability condition by which we may obtain eventual knowledge of the sign of the y -component.

Lemma 13. *Along any trajectory $\gamma(t)$, the quantity $Q = y/\omega$ is strictly decreasing. If $\gamma(t)$ satisfies (13), $\lim_{t \rightarrow T} Q = -\infty$. In particular, y eventually becomes negative.*

Proof. We compute

$$\frac{d}{dt} Q = -n \frac{x^2}{\omega} - \lambda \omega < 0.$$

Integrating, we find that, for any $S < t_0 < t < T$,

$$\frac{y}{\omega}(t) \leq \frac{y}{\omega}(t_0) - \lambda \int_{t_0}^t \omega(\sigma) d\sigma.$$

□

The following observation is also immediate.

Lemma 14. *If $\gamma(t)$ satisfies (13), then $\limsup_{t \rightarrow T} x(t) \geq 0$.*

Proof. If $x(t) < -\delta$ on some $(a, T) \subset (S, T)$, then $\omega(t) \leq Ce^{-\delta t}$ on the same interval, and $\gamma(t)$ cannot satisfy (13). $\square$

Thus, in view of Lemma 11, no trajectory satisfying (13) can enter the region $\{x < -1\}$. That no trajectory satisfying (13) can enter the region $\{x > 1\}$ is true (as we prove next), but less obvious since $\omega(t) \rightarrow \infty$. Taken together, Lemmas 11, 14, and 15 prove that any complete metric on $\mathbb{R}^{n+1}$ satisfies $\nu_2 \geq 0$.

The proof follows the lines of an argument due to Bryant and Ivey (c.f. [RF2]) demonstrating the incompleteness of a similar trajectory of the steady soliton system. In fact, if one regards the trajectories of (9) in the plane $\{\omega = 0\}$ as trajectories of the analogous system for steady solitons (cf. the remarks at the end of Section 2.3), then Ivey's argument pertains to the trajectory in the plane emerging from P_0 in the direction opposite the Bryant soliton. The following lemma then may be viewed as an extension of his finding to the neighboring family of trajectories in the unstable manifold U_0 which populate the sector between the flat trajectory with $x \equiv 0$ and the plane $\{\omega = 0\}$. These trajectories correspond to metrics of strictly negative sectional curvature and are all incomplete. However, Ivey's argument does not carry over directly, as, in the expression for $\frac{dx}{dt}$ in the shrinking case, one has to contend with an additional term $(-\lambda\omega^2)$ of uncooperative sign.

Lemma 15. *Suppose $x(t_0) > 1$ and $\frac{dx}{dt}(t_0) > 0$ at some $t_0 \in (S, T)$. Then $\int_{t_0}^T \omega(\sigma) d\sigma < \infty$.*

Proof. We proceed by contradiction. Suppose $\gamma(t)$ satisfies (13). Then, by (11), $x(t) > 1$ and $\frac{dx}{dt}(t) > 0$ for $t > t_0$, and, by Lemma 13, there is a $t_1 \in (t_0, T)$ such that $y(t) < 0$ for all $t \geq t_1$.

Hence, by (11), we have

$$(15) \quad \frac{d^2x}{dt^2} \geq (n-1)x(x^2-1) + 3x\frac{dx}{dt} > \frac{3}{2}\frac{d(x^2)}{dt}.$$

Now, since the interval (S, T) is assumed maximal, and since bounds on x imply bounds on the derivatives of y and ω , if $T < \infty$ we must have $\limsup_{t \rightarrow T} |x(t)| = \lim_{t \rightarrow T} x = \infty$. On the other hand, $x(t)$ is uniformly convex by (15), so even if $T = \infty$ we still have $\lim_{t \rightarrow T} x(t) = \infty$. Returning to (15) with this fact in hand, we find

$$\frac{dx}{dt} \geq \frac{5}{4}x^2 + 1$$

for all t greater than some $t_2 \geq t_1$. (The coefficient $5/4$ is chosen for convenience and could be replaced by $3/2 - \epsilon$ for any $\epsilon > 0$ – below, we merely require it to be greater than one.) From this equation it follows that $T < \infty$ and

$$\frac{d}{dt} \arctan \left(\frac{\sqrt{5}}{2}x \right) \geq \frac{\sqrt{5}}{2},$$

which implies

$$x(t) \leq \frac{4}{5(T-t)}$$

for t sufficiently close to T . Since $\frac{d}{dt} \log \omega = x$, integrating and applying the above bound yields

$$\omega(t) \leq \frac{C_2}{(T-t)^{\frac{4}{5}}}$$

for some constant C_2 , contradicting (13). $\square$

Together, Lemmas 10 and 15 allow us to restrict our attention to trajectories which remain in the region $\{-1 < x < 1\}$, and, consequently, to those with infinite existence time $t \in (S, \infty)$ since a trajectory with x bounded cannot satisfy condition (13) on a interval bounded above. Along such trajectories, y becomes negative and Lemma 11 implies that eventually x acquires a constant sign. As a consequence, we obtain the following refinement of Lemma 13:

Lemma 16. *If $\gamma(t)$ is a trajectory of (9) defined on (S, ∞) satisfying condition (13) and $-1 < x(t) < 1$, then $\lim_{t \rightarrow \infty} y(t) = -\infty$.*

Proof. Suppose $\limsup_{t \rightarrow \infty} y(t) \geq -M$ for some $M > 1$, and choose $t_k \in (S, \infty)$ such that $t_k \nearrow \infty$ and $y(t_k) \geq -M$. Then, by Lemma 13, $\lim_{k \rightarrow \infty} \omega(t_k) = 0$. Since x eventually acquires a constant sign, and $\frac{d}{dt}\omega = x\omega$, we must have $x(t) \leq 0$ eventually, and $\omega(t)$ must tend to 0 outright as $t \rightarrow \infty$.

Now, equation (11) shows that $x(t)$ cannot attain a local maximum on $\{-1 < x \leq 0\}$ unless $x = 0$, and we know $x = 0$ and $\frac{dx}{dt} = 0$ simultaneously only if $x \equiv 0$, in which case $dy = -(n-1)dt$. Otherwise, x is eventually monotonic in t and either increases or decreases to a limit $\bar{x} \in [-1, 0]$. By the remarks preceding this lemma, we cannot have $\bar{x} < 0$ if the trajectory is to satisfy condition (13). So assume $\bar{x} = 0$, which implies $\frac{dx}{dt} > 0$ eventually. Then, for any ϵ , we can choose t_ϵ such that $t > t_\epsilon$ implies both $y(t) < 0$ and $nx^2 + \lambda\omega^2 < \epsilon$. For such t , we have

$$\frac{dy}{dt} = xy - nx^2 - \lambda\omega^2 > -\epsilon.$$

Fixing $\epsilon < \frac{n-1}{2}$, we find, each for k ,

$$y(t) > -M - \epsilon(t - t_k)$$

and

$$\frac{dx}{dt}(t) = x^2 - xy + n - 1 - \lambda\omega^2 > \frac{n-1}{2} + x(t_k)(M + \epsilon(t - t_k)),$$

(where, in obtaining the last inequality, we used that x is monotonically increasing).

Hence,

$$x(t) - x(t_k) > \left(\frac{n-1}{2} + Mx(t_k) \right) (t - t_k) + \epsilon \frac{x(t_k)}{2} (t - t_k)^2.$$

For $k \gg 0$, $Mx(t_k) > -(n-1)/4$, so that the above (with the monotonicity of $x(t)$) implies that there exists $\delta = \delta(M, \epsilon, n) > 0$ and a subsequence $t_{k_j} \rightarrow \infty$ such that $x(t_{k_{j+1}}) > x(t_{k_j}) + \delta$ for all j . This contradicts that $x \nearrow 0$, and proves $\limsup_{t \rightarrow \infty} y(t) = -\infty$. $\square$

3.4. Proof of the case $\mathcal{M}^{n+1} \approx \mathbb{R}^{n+1}$. The results of the last section are enough to assemble the

Proof of Claim (2) of Theorem 1. Since the underlying manifold $\mathcal{M}^{n+1}$ is diffeomorphic to $\mathbb{R}^{n+1}$, the smooth extension of the metric to the origin $r = 0$ requires $S = -\infty$ and our solution $\gamma(t) = (\omega(t), y(t), \omega(t))$ of (9) to satisfy $\lim_{t \rightarrow -\infty} \gamma(t) = P_0 = (0, 1, n)$.

We claim first that if our trajectory is to satisfy condition (13), then $x \leq 1$. For if ever $x > 1$, since $\lim_{t \rightarrow -\infty} x(t) = 1$, we would have to have $\frac{dx}{dt}(t_0) > 0$ and

$x(t_0) > 1$ at some earlier t_0 . But then, by Lemmas 10 and 15, the x -component would blow-up too fast for $\gamma(t)$ to satisfy (13). So we must have $x \leq 1$ for all t .

Then, if ever $x = 1$, we must also have $\frac{dx}{dt} = 0$ at the same time, which, as pointed out in Remark 8, happens only if $x \equiv 1$ —i.e., only if $\gamma(t)$ corresponds to the flat solution of Example 6. We claim that this is the only trajectory emanating from P_0 which satisfies (13).

We may now assume that $x(t) < 1$ on our trajectory and that for some t_0 (hence all $t < t_0$), $\frac{dx}{dt}(t_0) < 0$ and $0 < x(t_0) < 1$. Since, by equation (11), $\frac{d^2x}{dt^2}$ is strictly negative at all critical points of x in the region $0 < x < 1$, there are two possibilities for our trajectory: either

- (1) x decreases monotonically to a limit $\bar{x} \in [0, 1)$ as $t \rightarrow \infty$, or
- (2) $\gamma(t)$ enters the region $\{x \leq 0\}$ at some time $t = t_1$.

Knowing that $y \rightarrow -\infty$ as $t \rightarrow \infty$ (in fact, just knowing that eventually $y < 0$ suffices), we can dispose of case (1) by observing that while $x \in [0, 1]$,

$$(16) \quad \frac{d^2x}{dt^2} \leq (n-1)x(x^2-1) + \epsilon \frac{dx}{dt} < 0$$

once $y < -\epsilon < 0$. Hence x eventually becomes negative.

Now, we also know from Lemma 14 that x cannot tend to a negative limit or become strictly less than -1 if condition (13) is to be satisfied. Since $\frac{d^2x}{dt^2}$ is strictly positive at critical points of x in the region $\{-1 < x < 0\}$, and since the only trajectories with critical points of x on the boundary of this region are classified in Examples 6 and 7 and neither emanate from P_0 , we face only two alternatives:

- (2a) either γ enters the region $x > 0$ again, or
- (2b) $x \nearrow 0$ as $t \rightarrow \infty$.

Alternative (2a) is immediately excluded by Lemmas 11 and 15: no trajectory which emanates from P_0 can satisfy $x(t_0) = 0$, $\frac{dx}{dt}(t_0) > 0$, $y(t_0) \geq 0$, and no trajectory which satisfies $x(t_0) = 0$, $\frac{dx}{dt}(t_0) > 0$, $y(t_0) \leq 0$ can satisfy (13).

For (2b), we observe that since $y \rightarrow -\infty$ as $t \rightarrow \infty$, we have $3x - y > \epsilon > 0$ eventually, and thus we may obtain the analog of equation (16) for t sufficiently large

$$(17) \quad \frac{d^2x}{dt^2} = (n-1)x(x^2-1) + (3x-y)\frac{dx}{dt} > \epsilon \frac{dx}{dt} > 0,$$

which is incompatible with $x \nearrow 0$ as $t \rightarrow \infty$.

The trajectory $x \equiv 1$ is therefore the unique trajectory emanating from P_0 satisfying (13), and the proof of the case $\mathcal{M}^{n+1} \approx \mathbb{R}^{n+1}$ is complete. $\square$

3.5. Proof of the case $\mathcal{M}^{n+1} \approx \mathbb{R} \times S^n$. As remarked earlier, the results in Section 3.3, regarding trajectories satisfying the forwards extendability condition (13) have natural analogs for trajectories satisfying the backwards version (13).

Lemma 17.

- (1) Suppose $\gamma(t)$ satisfies (14).
 - (a) $\lim_{t \rightarrow S} Q(t) = \infty$ and y is initially positive.
 - (b) $\liminf_{t \rightarrow S} x(t) \leq 0$
 - (c) If $-1 < x(t) < 1$ for all t (so $-S = T = \infty$), then $\lim_{t \rightarrow -\infty} y(t) = \infty$.

(2) If $x(t_0) < -1$ and $\frac{dx}{dt}(t_0) > 0$ at some $t_0 \in (S, T)$,

$$(18) \quad \int_S^{t_0} \omega(\sigma) d\sigma < \infty$$

Proof. Let $\bar{\gamma}(t) = L(\gamma(\tau(t)))$ be as in Remark 4.

$$(19) \quad \int_S^{t_0} \omega(\sigma) d\sigma = \int_{t_0}^{\bar{T}} \bar{\omega}(\sigma) d\sigma.$$

Thus, if $\gamma(t)$ satisfies (14), $\bar{\gamma}(t)$ satisfies (13) and the claims of part 1 follow by the application of Lemmas 13, 14, and 16 to $\bar{\gamma}(t)$.

For part 2, note that $\bar{x}(t_0) > 1$, $\frac{d\bar{x}}{dt}(t_0) > 0$ if $x(t_0) < -1$, $\frac{dx}{dt}(t_0) > 0$, so Lemma 15 and equation (19) yield the inequality (18). $\square$

Now we turn to the remainder of the proof of Theorem 1.

Proof of Claim 3 of Theorem 1. We shall show that the only trajectory $\gamma(t)$ satisfying both (13) and (14) is that of Example 7 with $x \equiv 0$ and $\omega \equiv \sqrt{(n-1)/\lambda}$.

By Lemmas 10 and 15 and case (2) of Lemma 17, we can assume that $-1 \leq x(t) \leq 1$ and consequently also $S = -\infty$, $T = \infty$. In fact, since neither of the trajectories with $x \equiv \pm 1$ can satisfy both conditions (13) and (14), we may assume $-1 < x(t) < 1$.

By Lemma 14 and part (1b) of Lemma 17, we know $\liminf_{t \rightarrow -\infty} x(t) \leq 0$ and $\limsup_{t \rightarrow \infty} x(t) \geq 0$. Thus since $\frac{d^2x}{dt^2}$ is strictly positive at critical points of x in the region $\{-1 < x < 0\}$ and strictly negative at these critical points in the region $\{0 < x < 1\}$, either $\gamma(t)$ crosses the plane $\{x = 0\}$ at some time or its x -component maintains a constant sign and tends monotonically to 0 as $t \rightarrow \infty$ or $t \rightarrow -\infty$. We claim that this latter option cannot occur. For, in light of the remarks in the preceding paragraph, the only scenarios of this option in which $\gamma(t)$ could potentially satisfy the extendability criteria would be $x < 0$ and $x \nearrow 0$ as $t \rightarrow \infty$, or $x > 0$ and $x \searrow 0$ as $t \rightarrow -\infty$. But the case $x < 0$, $x \nearrow 0$ was eliminated in the argument for Claim (2), and the case $x \searrow 0$ as $t \rightarrow -\infty$ reduces to the previous one by the consideration of the trajectory $\bar{\gamma}(t)$.

Thus, we conclude that there must exist a t_0 such that $x(t_0) = 0$. If $\frac{dx}{dt}(t_0) \neq 0$, then we may assume $\frac{dx}{dt}(t_0) > 0$, as the argument given in the case $\mathcal{M}^{n+1} \approx \mathbb{R}^{n+1}$ implies that the only trajectories satisfying (13) and $\frac{dx}{dt}(t_0) < 0$ initially lie in the region $\{x < -1\}$ and cannot thus satisfy (14). However, if $\frac{dx}{dt}(t_0) > 0$, then Lemma 11 implies that either again $\gamma(t)$ lies initially in the region $\{x < -1\}$ or eventually in the region $\{x > 1\}$, in which case, by Lemma 15 and part (2) of Lemma 17, $\gamma(t)$ can satisfy at most one of the conditions (13) and (14).

Thus we can only have $\frac{dx}{dt}(t_0) = 0$, which implies that γ coincides with the trajectory $x \equiv 0$, $\omega \equiv \sqrt{(n-1)/\lambda}$ as claimed. $\square$

Acknowledgement. *The author wishes to thank Professors Bennett Chow and Lei Ni for their support, encouragement, and for useful discussions pertaining to this work. The author is especially indebted to Professor Thomas Ivey, whose analysis of the problem in the expanding and steady soliton cases served as a guide for the study of the shrinking soliton case presented here.*

REFERENCES

- [BW] C. Böhm and B. Wilking, *Manifolds with positive curvature are space forms* preprint.
- [CK] B. Chow and D. Knopf, *The Ricci Flow: An Introduction*. AMS, Providence, RI, (2004).
- [CLN] B. Chow, P. Lu, and L. Ni, *Hamilton's Ricci flow*. Lectures in Contemporary Mathematics, Science Press and Graduate Studies in Mathematics, American Mathematical Society (co-publication). To appear.
- [CLT] X. Chen, P. Lu, and G. Tian, *A note on uniformization of Riemann surfaces by Ricci flow*. arXiv:math.DG/0505163 v2.
- [FIK] M. Feldman, T. Ilmanen, D. Knopf. *Rotationally symmetric shrinking and expanding gradient Kähler-Ricci solitons*. J. Diff. Geom. **65** (2003), no. 2, 169–209.
- [H] R. Hamilton, *The formation of singularities in the Ricci flow*, Surveys in Differential Geometry **2** (1995) pp. 7-136, International Press.
- [H2] R. Hamilton, *The Ricci flow on surfaces*, Contemp. Math. **71** AMS, Providence, RI, (1988), 237-262.
- [H4] R. Hamilton, *Four-manifolds with positive curvature operator*, J. Differential Geom., **24** (1986) pp. 153-179.
- [I] T. Ivey, *On solitons for the Ricci flow*. Ph.D. Thesis, Duke University, (1992).
- [P2] G. Perelman. *Ricci flow with surgery on three-manifolds* arXiv:math.DG/0303109
- [RF2] B. Chow, et al. *The Ricci Flow: Techniques and Applications*. AMS, Providence, RI. (In preparation)

UNIVERSITY OF CALIFORNIA, SAN DIEGO
E-mail address: bkotschw@math.ucsd.edu